

On Frattini arguments in L -groups of finite Morley rank

Jeffrey Burdges*

April 20, 2009

Proposition 0.1. *Let $\hat{G}$ be a group of finite Morley rank, let G itself be a definable connected normal subgroup group of $\hat{G}$, and let S be a Sylow 2-subgroup of $\hat{G}$. Then G has an S -invariant Carter subgroup.*

We use the following facts, as well as the Frécon-Jaligot construction and the conjugacy of descent tori.

Fact 0.2 ([Bur06, Cor. 3.5]). *Let H be a nilpotent group of finite Morley rank. Then $H = D*B$ is a central product of definable characteristic subgroups $D, B \leq H$ where D is divisible and B has bounded exponent (which is connected iff H is connected). Let T be the torsion part of D . Then we have decompositions of D and B as follows.*

$$\begin{aligned} D &= d(T) * U_{0,1}(H) * U_{0,2}(H) * \cdots \\ B &= U_2(H) \oplus U_3(H) \oplus U_5(H) \oplus \cdots \end{aligned}$$

Fact 0.3 ([Bur06, Lemma 4.4]). *Let H be a connected solvable group of finite Morley rank. Suppose that S is a nilpotent $U_{0,r}$ -subgroup of H , and that $H = U_p(H)S$ for some p prime. Then H is nilpotent, and $[U_p(H), S] = 1$.*

Fact 0.4 ([Bur06, Lemma 4.4]). *Let $H = KT$ be a group of finite Morley rank with $K \triangleleft H$ a nilpotent $U_{0,r}$ -group and T a nilpotent $U_{0,s}$ -group for some $s \geq r$. Then H is nilpotent.*

Proof of Proposition 0.1. We take G to be a minimal counterexample, i.e. proper connected S -invariant subgroups of G have S -invariant Carter subgroups. Then G is nonsolvable by a Frattini argument. We observe that Carter subgroups of G correspond to Carter subgroups of $G/Z^\circ(G)$. So G is centerless too. We may also assume that $C_S(G) = 1$.

If G contains divisible torsion, we take T to be a maximal descent torus of G (exists by [Che05]). By [Che05], all maximal descent tori are conjugate, so $G =$

*Supported by NSF grant DMS-0100794, and Deutsche Forschungsgemeinschaft grant Te 242/3-1.

$N_G(T)G$ by a Frattini argument. Hence a conjugate of T is S -invariant. There is a Carter subgroup Q containing T , by a Frécon-Jaligot construction. By Fact [BN94, Thm. 6.16], T is central in $M := N_G^\circ(T)$. Now we have $M < G$, since G is centerless. Since M is S -invariant, there is an S -invariant Carter subgroup Q of M . Since T is a characteristic subgroup of Q , $N_G^\circ(Q) \leq N_G^\circ(T) \leq M$. So Q is a Carter subgroup of G . Hence we may assume that G has no divisible torsion.

We find a maximal S -invariant group R with “minimal degree of unipotence” r . Let r be the minimal reduced rank such that there is an S -invariant nilpotent $U_{0,r}$ -subgroup of G , or take $r = \infty$ if there are no S -invariant nilpotent $U_{0,r}$ -subgroups. Let i be a central involution of S . Since G is nonabelian, $C_G^\circ(i)$ is infinite by [BN94, Ex. 13 & 15 on p. 79]. and clearly S -invariant. Since $C_G^\circ(i) < G$, $C_G^\circ(i)$ has a definable S -invariant nilpotent subgroup, by induction. So, if $r = \infty$, there is an S -invariant nilpotent group of bounded exponent, by Fact 0.2. Let R be a maximal S -invariant nilpotent $U_{0,r}$ -subgroup of G , or else a maximal S -invariant p -unipotent subgroup of G if $r = \infty$. Now consider the S -invariant group $M := N_G^\circ(R)$.

First suppose that $M < G$. Then there is an S -invariant Carter subgroup Q of M , by induction. Now RQ is solvable. We show that R is characteristic in Q . If $r = \infty$, RQ is nilpotent by Fact 0.3, and RQ has bounded exponent. So $R = Q$ by maximality. If $r < \infty$, r is the minimal reduced rank found in Q , since $U_{0,t}(Q)$ is S -invariant for all t . So RQ is nilpotent by Fact 0.4. $R \leq Q$, and hence $U_{0,r}(Q) = R$. In either case, $N_G^\circ(Q) \leq M$. So Q is a Carter subgroup of G .

Next suppose that $M = G$, i.e. $R \triangleleft G$. Then there is an S -invariant Carter subgroup $\bar{Q}$ of $\bar{G} = G/R$. Now the pullback Q of $\bar{Q}$ to G is solvable, and $R \leq Q$. Since G is nonsolvable, $\bar{Q} < \bar{G}$, and $Q < G$. Since Q is S -invariant, there is an S -invariant Carter subgroup L of Q , by induction. We show that $Q = RL$ is nilpotent. If $r = \infty$, RL is nilpotent by Fact 0.3. If $r < \infty$, r is the minimal reduced rank found in L (or Q), since $U_{0,t}(L)$ is S -invariant for all t . So RL is nilpotent by Fact 0.4, and $R \leq L$. By maximality of R , $U_{0,r}(L) = R$. By Fact 0.2. $U_{0,t}(L)$ centralizes R for $t > r$. In either case, $Q = RL$ is nilpotent. We observe that $N_G(Q)/R \leq N_{\bar{G}}(\bar{Q})$ has the same rank as $\bar{Q}$. So $\text{rk}(N_G(Q)) = \text{rk}(Q)$, and $N_G^\circ(Q) = Q$. Thus Q is an S -invariant Carter subgroup of G . $\square$

References

- [BN94] Alexandre Borovik and Ali Nesin. *Groups of Finite Morley Rank*. The Clarendon Press Oxford University Press, New York, 1994. Oxford Science Publications.
- [Bur06] Jeffrey Burdges. Sylow 0-unipotent subgroups in groups of finite morley rank. *J. Group Theory*, 2006. To appear.
- [Che05] Gregory Cherlin. Good tori in groups of finite Morley rank. *J. Group Theory*, 8:613–621, 2005.